

## 兔抗 TP53I3 多克隆抗体

中文名称: 兔抗 TP53I3 多克隆抗体

英文名称: Anti-TP53I3 rabbit polyclonal antibody

别 名: PIG3

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

抗 原: TP53I3

宿 主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: Unconjugate

## 技术规格

**Background:** 

The protein encoded by this gene is similar to oxidore ductases, which are enzymes involved in cellular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppressor p53 and is thought to be involved in p53-mediated cell death. It contains a p53 consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucleotide microsatellite sequence. The microsatellite is polymor phic, with a varying number of pentanucleotide repeats



	directly correlated with the extent of transcriptional act ivation by p53. It has been suggested that the microsa tellite polymorphism may be associated with differentia I susceptibility to cancer. Alternatively spliced transcript variants encoding different isoforms have been found f or this gene.
Applications:	WB
Name of antibody:	TP53I3
Immunogen:	Synthesized peptide derived from internal of human Q ORX.
Full name:	tumor protein p53 inducible protein 3
Synonyms:	PIG3
SwissProt:	Q53FA7
WB Predicted band size:	36 kDa
WB Positive control:	293 cells lysate
WB Recommended dilution:	500-3000

