

## 兔抗 IL17D 多克隆抗体

中文名称：兔抗 IL17D 多克隆抗体

英文名称：Anti-IL17D rabbit polyclonal antibody

别名：IL27; IL-27; IL-17D

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：IL17D

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>WB Recommended dilution:</b>	200-1000
<b>WB Positive control:</b>	Human placenta tissue and A431 cell
<b>WB Predicted band size:</b>	22 kDa
<b>IHC Recommend dilution:</b>	50-200
<b>IHC positive control:</b>	Human tonsil and human gastric cancer
<b>ELISA Recommended dilution:</b>	5000-10000
<b>SwissProt:</b>	Q8TAD2
<b>Synonyms :</b>	IL27; IL-27; IL-17D

<b>Full name:</b>	interleukin 17D
<b>Immunogen:</b>	Synthetic peptide of human IL17D
<b>Name of antibody:</b>	IL17D
<b>Applications:</b>	ELISA, WB, IHC
<b>Background:</b>	The protein encoded by this gene is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines including IL6, IL8 and CSF2/ GM-CSF. The increased expression of IL8 induced by this cytokine was found to be NF-kappa B-dependent.



