

## 兔抗 DALRD3 多克隆抗体

中文名称：兔抗 DALRD3 多克隆抗体

英文名称：Anti-DALRD3 rabbit polyclonal antibody

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：DALRD3

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | The exact function of this gene is not known. It encodes a protein with a DALR anticodon binding domain similar to that of class Ia aminoacyl tRNA synthetases. This gene is located in a cluster of genes (with a complex sense-anti-sense genome architecture) on chromosome 3, and contains two micro RNA (miRNA) precursors (mir-425 and mir-191) in one of its introns. Preferential expression of this gene (the miRNAs and other genes in the cluster) in testis suggests a role of this gene in spermatogenesis (PMID:19906709). |
| <b>Applications:</b>     | ELISA, WB, IHC   |
| <b>Name of antibody:</b> | DALRD3   |
| <b>Immunogen:</b>        | Fusion protein of human DALRD3   |
| <b>Full name:</b>        | DALR anticodon binding domain containing 3   |
| <b>SwissProt:</b>        | Q5D0E6   |

|                                    |                    |
|------------------------------------|--------------------|
| <b>ELISA Recommended dilution:</b> | 5000-10000         |
| <b>IHC positive control:</b>       | Human liver cancer |
| <b>IHC Recommend dilution:</b>     | 25-100             |
| <b>WB Predicted band size:</b>     | 59 kDa             |
| <b>WB Positive control:</b>        | HepG2 cell lysate  |
| <b>WB Recommended dilution:</b>    | 500-2000           |



