

兔抗 EIF4E2 多克隆抗体

中文名称：兔抗 EIF4E2 多克隆抗体

英文名称： Anti-EIF4E2 rabbit polyclonal antibody

别名： eukaryotic translation initiation factor 4E family member 2; 4EHP; IF4e; 4E-LP; h4EHP; EIF4EL3

相关类别： 一抗

储存： 冷冻（-20℃）

宿主： Rabbit

抗原： EIF4E2

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:	Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation (PubMed:9582349, PubMed:17368478, PubMed:25624349). Acts as a repressor of translation initiation (PubMed:22751931). In contrast to EIF4E, it is unable to bind eIF4G (EIF4G1, EIF4G2 or EIF4G3), suggesting that it acts by competing with EIF4E and block assembly of eIF4F at the cap (By similarity).
Applications:	ELISA, IHC
Name of antibody:	EIF4E2
Immunogen:	Fusion protein of human EIF4E2

Full name:	eukaryotic translation initiation factor 4E family member 2
Synonyms:	4EHP; IF4e; 4E-LP; h4EHP; EIF4EL3
SwissProt:	O60573
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	50-200

