

## 兔抗 RACGAP1(Phospho-Ser387) 多克隆抗体

中文名称: 兔抗 RACGAP1(Phospho-Ser387) 多克隆抗体

英文名称: Anti-RACGAP1(Phospho-Ser387) rabbit polyclonal antibody

别名: CYK4; ID-GAP; HsCYK-4; MgcRacGAP

相关类别: 一抗

储存: 冷冻 (-20℃) 避光

宿主: Rabbit

抗原: RACGAP1(Phospho-Ser387)

反应种属: Human Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

### 技术规格

|                          |  |
|--------------------------|--|
| <b>Background:</b>       | The protein encoded by this gene belongs to the GTPase-activating protein (GAP) family. GAPs bind activated forms of Rho GTPases and stimulate GTP hydrolysis. Through this catalytic function, GAPs negatively regulate Rho-mediated signals. This protein plays a regulatory role in initiation of cytokinesis, controlling cell growth and differentiation of hematopoietic cells, regulating spermatogenesis, and in neuronal proliferation. |
| <b>Applications:</b>     | WB   |
| <b>Name of antibody:</b> | RACGAP1(Phospho-Ser387)  |
| <b>Immunogen:</b>        | Peptide sequence around phosphorylation site of serine 387 (R-I-S(p)-G-C) derived from Human GT  |

|                                 |                                  |
|---------------------------------|----------------------------------|
|                                 | Pase Activating Protein.         |
| <b>Full name:</b>               | Rac GTPase activating protein 1  |
| <b>Synonyms :</b>               | CYK4; ID-GAP; HsCYK-4; MgcRacGAP |
| <b>SwissProt:</b>               | Q9H0H5                           |
| <b>WB Predicted band size:</b>  | 71 kDa                           |
| <b>WB Positive control:</b>     | COS7 cells lysates               |
| <b>WB Recommended dilution:</b> | 500-1000                         |

