

兔抗 PLD2(Phospho-Tyr169) 多克隆抗体

中文名称:	兔抗 PLD2(Phospho-Tyr169) 多克隆抗体
英文名称:	Anti-PLD2(Phospho-Tyr169) rabbit polyclonal antibody
别名:	PLD2
相关类别:	一抗
储存:	冷冻 (-20℃) 避光
宿主:	Rabbit
抗原:	PLD2(Phospho-Tyr169)
反应种属:	Human
标记物:	Unconjugate
克隆类型:	rabbit polyclonal

技术规格

Background:	The protein encoded by this gene catalyzes the hydrolysis of phosphatidylcholine to phosphatidic acid and choline. The activity of the encoded enzyme is enhanced by phosphatidylinositol 4,5-bisphosphate and ADP-ribosylation factor-1. This protein localizes to the peripheral membrane and may be involved in cytoskeletal organization, cell cycle control, transcriptional regulation, and/or regulated secretion. Two transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2011]
Applications:	WB, IHC
Name of antibody:	PLD2(Phospho-Tyr169)
Immunogen:	Peptide sequence around phosphorylation site of tyrosine 169(E-N-Y(p)-L-N) derived from Human PLD2.
Full name:	phospholipase D2

Synonyms :	PLD2
SwissProt:	O14939
IHC positive control:	Human brain tissue
IHC Recommend dilution:	50-100
WB Predicted band size:	106 kDa
WB Positive control:	Jurkat cells lysates treated with TNF
WB Recommended dilution:	500-1000



