

## 兔抗 PPARA 多克隆抗体

中文名称: 兔抗 PPARA 多克隆抗体

英文名称: Anti-PPARA rabbit polyclonal antibody

别 名: PPAR, NR1C1, hPPAR, PPARalpha

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗 原: PPARA

反应种属: Human, Mouse, Rat

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

**Background:** 

Peroxisome proliferators include hypolipidemic drugs, her bicides, leukotriene antagonists, and plasticizers; this term arises because they induce an increase in the size and n umber of peroxisomes. Peroxisomes are subcellular organ elles found in plants and animals that contain enzymes f or respiration and for cholesterol and lipid metabolism. T he action of peroxisome proliferators is thought to be m ediated via specific receptors, called PPARs, which belong to the steroid hormone receptor superfamily. PPARs affect the expression of target genes involved in cell proliferation, cell differentiation and in immune and inflammation responses. Three closely related subtypes (alpha, beta/del ta, and gamma) have been identified. This gene encodes the subtype PPAR-alpha, which is a nuclear transcription



## 全国订货电话 4008-723-722

	factor. Multiple alternatively spliced transcript variants hav e been described for this gene, although the full-length nature of only two has been determined.
Applications:	ELISA, WB, IHC
Name of antibody:	PPARA
Immunogen:	Synthetic peptide of human PPARA
Full name:	peroxisome proliferator activated receptor alpha
Synonyms:	PPAR; NR1C1; hPPAR; PPARalpha
SwissProt:	Q07869
<b>ELISA Recommended dilution:</b>	5000-10000
IHC positive control:	Human lung cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	52 kDa
WB Positive control:	Rat heart tissue lysate
WB Recommended dilution:	200-1000





