

兔抗 PRKCD (Phospho-Ser645)多克隆抗体

中文名称:	兔抗 PRKCD (Phospho-Ser645)多克隆抗体
英文名称:	Anti-PRKCD (Phospho-Ser645) rabbit polyclonal antibody
别名:	MAY1; PKCD; nPKC-delta
相关类别:	一抗
储存:	冷冻 (-20℃) 避光
宿主:	Rabbit
抗原:	PRKCD (Phospho-Ser645)
反应种属:	Human Rat
标记物:	Unconjugate
克隆类型:	rabbit polyclonal

技术规格

Background:	This is calcium-independent, phospholipid-dependent, serine - and threonine-specific enzyme. PKC is activated by diacyl glycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters. May play a role in antigen-dependent control of B-cell function. Phosphorylates MUC1 in the C-terminal and regulates the interaction between MUC1 and beta-catenin.
Applications:	WB, IHC
Name of antibody:	PRKCD (Phospho-Ser645)
Immunogen:	Synthetic peptide of human PRKCD (Phospho-Ser645)
Full name:	protein kinase C, delta (Phospho-Ser645)
Synonyms :	MAY1; PKCD; nPKC-delta
SwissProt:	Q05655

IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	78 kDa
WB Positive control:	PC12 cells treated with PMA
WB Recommended dilution:	500-1000



