

兔抗 TGFBI 多克隆抗体

中文名称：兔抗 TGFBI 多克隆抗体

英文名称： Anti-TGFBI rabbit polyclonal antibody

别名： transforming growth factor beta induced; CSD; CDB1; CDG2; CSD1; CSD2; CSD3; EBMD; LCD1; BIGH3; CDGG1

相关类别： 一抗

储存： 冷冻 (-20℃)

宿主： Rabbit

抗原： TGFBI

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

Background:

This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondral bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy.

Applications:	ELISA, WB, IHC
Name of antibody:	TGFBI
Immunogen:	Fusion protein of human TGFBI
Full name:	transforming growth factor beta induced
Synonyms:	CSD; CDB1; CDG2; CSD1; CSD2; CSD3; EBMD; LCD1; BI GH3; CDGG1
SwissProt:	Q15582
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human liver cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	75 kDa
WB Positive control:	Rat liver tissue lysate
WB Recommended dilution:	500-2000



